



e-ISSN: 2456-6632

This content is available online at AESA

Archives of Agriculture and Environmental Science

Journal homepage: www.aesacademy.org



ORIGINAL RESEARCH ARTICLE



Effect of cassava peels and palm kernel cake (PKC) on the hematological and serum biochemical of grower pigs in Nigeria

H. Moseri², O.S. Oboh¹, S.E. Okosun¹ and C.E. Isidahomen^{1*}

¹Department of Animal Scimbrose Alli University Ekpoma, Edo State, NIGERIA

²Department of Agricultural Education, College of Education Agbor, Delta State, NIGERIA

*Corresponding author's E-mail: chukwukaraymond@gmail.com

ARTICLE HISTORY

Received: 14 September 2018
Revised received: 20 October 2018
Accepted: 10 November 2018

Keywords

Cassava (*Manihot esculenta*) peels
Hematological parameters
Palm kernel cake

ABSTRACT

The research was carried out for a period of ten weeks at the piggery unit of the teaching and research farm of the Faculty of Agriculture, Ambrose Alli University, Ekpoma, Edo State, Nigeria. The objective of this study was to assess the effect of cassava peels and palm kernel cake (PKC) at 50:50 ratio on the hematological and serum biochemical of grower pigs. Forty (40) grower pigs of large white were used; pigs were divided into 5 groups based on average initial weights (20-25kg) and each group of grower pigs were respectively allocated to each of the five treatment diets in a completely randomized design (CRD). At the end of the ten (10) weeks feeding trial, the animals were starved of feed for 12 hours before blood samples were collected from two pigs each per treatment from the ear vein using a sterilized disposable syringe and needle. The parameters determined were red blood cells (RBC), hemoglobin (Hb), packed cell volume (PCV), white blood cell (WBC), total protein, albumin, and globulin and serum cholesterol. The result indicated that hematological and serum biochemical parameters assessed fall within the normal range and is an indication of good nutrition and adequate nutrient intake. Therefore, the animals did not suffer muscular wastage but efficient utilization of cassava peels and palm kernel cake, ultimately resulted in high tissue deposition across treatment groups.

©2018 Agriculture and Environmental Science Academy

Citation of this article: Moseri, H., Oboh, O.S., Okosun, S.E. and Isidahomen, C.E. (2018). Effect of cassava peels and palm kernel cake (PKC) on the hematological and serum biochemical of grower pigs in Nigeria, *Archives of Agriculture and Environmental Science*, 3(4): 373-377, <https://dx.doi.org/10.26832/24566632.2018.030407>

INTRODUCTION

Cassava (*Manihot esculenta*) is widely grown in the tropical regions. It is estimated that most (60%) of the cassava crop is used for human consumption while the remainder (40%) is used for animal feed, starch and industrial applications (Oboh *et al.*, 2004). Since the early 1930s, cassava has been known and used as a livestock substitute instead of grains for monogastric animals. The popularity of cassava as a substitute resource in livestock feeds became fashionable first because of its relative cheapness compared to grains (especially corn) and later because of the increase demand for corn for human and industrial uses such as in textiles, breweries and bakeries (Moseri, 2010). Cassava roots can be fed to animals in different forms. It is advisable to have them processed before being included in the

pigs rations (Igene, 2006). Fresh roots of cassava are high in hydrogen cyanide which has been known to be poisonous to animals. Although the commercial production of dried cassava chip and pellets in most tropical countries (except Nigeria) is still insignificant, there is a considerable potential to expand the use of cassava in local animal feed markets and assessment of the effects on blood parameters of livestock. The blood volume in animals is important in order to understand and interpret packed cell volume, hemoglobin concentration, red blood cell counts, plasma protein concentrations, and other hematological values. According to Isaac *et al.* (2013), hematological components are valuable in measuring toxicity, especially with feed constituents that affect the blood as well as the physiological and health status of farm animals. Afolabi *et al.* (2010) posited that changes in hematological parameters are often used to

determine stresses due to nutrition and other factors. Hematological parameters are those parameters that are related to the blood and blood forming organs Waugh and Grant (2018); Bamishaiye et al. (2009) and are good indicators of the physiological status of animals Khan and Zarfar (2005). Hematological analysis involves the determination of different blood parameters which can be done using either electronic quantification or the manual quantification Etim et al. (2013). Hematological studies have been found useful for disease prognosis and for therapeutic as well as feed stress monitoring Togun et al. (2007) and Unigwe et al. (2016). Thus, the objective of this study is to evaluate the effect of cassava peels and palm kernel cake on hematological and serum biochemical of grower pigs.

MATERIALS AND METHODS

Experimental site

The research was carried out at the piggery unit of the teaching and research farm of the Faculty of Agriculture, Ambrose Alli University, Ekpoma. The farm is located in Esan West Local Government Area Council of Edo State, Nigeria; with an annual rainfall of 1500-2000mm per annual. Relative humidity is 75% and average temperature is 32°C.

Experimental animal and design

The research was carried out for a period of ten weeks. A total of 40 grower pigs of large white were used for the study. The pigs were divided into 5 groups based on average initial weight (20-25kg) and each group of grower pigs were respectively allocated to each of the five treatment diets in a completely randomized design (CRD). Each treatment group contained 2 replicates of 4 pigs (2male and 2female).

Management of animals

These pigs were fed twice daily and water supplied *ad libitum*. The treatment diets consisted of the following of cassava peels/Pkc at 0, 25, 50, 75 and 100% replacement of maize in the diet. The 50:50 ratio of cassava peels and palm kernel cake was derived by equal weighing (kg) of the two test ingredients percentage in the diet using a manual scale, all diets were formulated to be iso-nitrogenous and iso-caloric.

Hematological studies

At the end of the ten (10) weeks feeding phase, the animals were starved of feed for 12 hours before blood samples were collected from each pig from the ear vein using a sterilized disposable syringe and needle. Prior to bleeding, a cotton swab soaked in 70% ethanol was used to dilate the ear vein and to prevent infection or contamination of the blood sample. A 5.0ml blood was collected from each pig into labeled sterile universal bottles containing Ethylene-Diamine-Tetra-Acetic acid (EDTA) as anti-coagulant. This was used to determine the total red blood cells (RBC), hemoglobin (Hb), and packed cell volume (PCV), white blood cell (WBC). Another 5.0ml of blood was collected into labeled sterile sample bottles without anticoagulant and used to

determine the biochemical components viz., total protein, albumin, globulin and serum cholesterol according to AOAC (2000).

Statistical analysis

The data collected on various parameters were analyzed using FAO (2002) package. The difference in treatment means were separated using Duncan's New multiple range test as outlined by Obi (2002).

RESULTS AND DISCUSSION

With the exception of the neutrophils and MCHC, significant ($P<0.05$) differences were showed in blood parameters of grower pigs fed levels of cassava peels/Pkc experimental diets as presented in Table 1. Values for packed cell volume (PCV) showed significant ($P<0.05$) difference on the dietary treatments. Numerically, values were 44.50, 48.23, 48.45, 49.08, and 51.95(%) for treatments 0, 100, 50, 25 and 75(%) respectively. It was however indicated that treatment 0% was significantly ($P<0.05$) influenced compare to other treatments. Red blood cell (RBC) and hemoglobin (%) values were significantly ($P<0.05$) higher in treatment 4 (75%), with corresponding values of 8.13×10^6 /ml and 12.98g/dl, while the lowest was in control (0%) diet in both RBC and Hb with values of 7.17×10^6 /ml and 11.48g/dl, respectively. White blood cell (WBC) was lowest (26.45×10^3 /ml) among grower pigs fed 25% treatment diets and highest in treatment 1 (39.50×10^3 /ml). MCV value ranged from 59.90fl in diet 1 to 64.00 in diet 4. MCH was least in control (0%) with the value 15.55(pg) and higher value was obtained in 75% (17.25pg), respectively. Mean corpuscular hemoglobin concentration (MCHC) recorded values of 25.95, 26.30, 26.50, 26.90 and 26.52% for treatment 1, 2, 3, 4 and 5 respectively. Values recorded for neutrophils were 60.80, 59.73, 57.35, 55.45, and 51.60% for 1, 2, 3, 4 and 5.

Serum biochemical parameters of grower pigs fed experimental diets are shown in Table 2. Serum biochemical parameters for grower pigs fed dietary treatments were significantly ($P<0.05$) affected on the total protein, albumin, creatinine, urea and globulin for all treatment groups. Total protein values recorded ranged from 8.05(g/dl) for treatment 5 to 8.39(g/dl) in treatment 1. Albumin values of 3.50, 3.60, 3.40, 4.10 and 3.75g/dl for treatments 1, 2, 3, 4 and 5 respectively. Globulin recorded values of 4.89, 4.50, 4.80, 4.60 and 4.30g/dl for treatments 1, 2, 3, 4 and 5 respectively. Creatinine values were 1.00, 0.80, 1.10, 0.93 and 0.87g/dl while urea values of 40.50, 39.50, 28.50, 45.50 and 35.50g/dl for treatment 1, 2, 3, 4 and 5. Cholesterol values ranged from 109.00 to 118.25mg/dl from experimental diet 5 to 1, significant ($P<0.05$) differences were recorded for treatment 1, 2 and 3. However, no significant ($P>0.05$) differences observed between 4 and 5 with corresponding values of 118.25, 116.30, 114.00, 110.00 and 109.00mg/dl for treatment 1, 2, 3, 4 and 5, respectively.

The importance of blood cannot be over emphasized as we all know that blood plays a vital role in the life of all living

Table 1. Composition of experimental diet for growers pig.

Ingredients	0	25%	50%	75%	100%
Maize	45.00	33.75	22.50	11.25	---
Cassava peels/PKC	---	11.25	22.50	33.75	45.00
Ground Nut Cake	18.87	20.0	21.13	22.28	23.42
Wheat Offal	30.83	28.70	27.37	26.02	24.68
Bone Meal	1.50	1.50	1.50	1.50	1.50
Limestone	2.00	2.00	2.00	2.00	2.00
Palm Oil	1.00	2.00	2.20	2.40	2.60
Growers Premix*	0.25	0.25	0.25	0.25	0.25
Salt	0.35	0.35	0.35	0.35	0.35
Ronozyme**	0.20	0.20	0.20	0.20	0.20
Total	100.00	100.00	100.00	100.00	100.00
Calculated analysis:					
Crude Protein (%)	17.00	17.00	17.00	17.00	17.00
ME (Keal/Kg)	2824	2804	2783	2761	2739
Fibre (%)	5.07	6.72	8.37	10.01	11.64
Ash (%)	5.88	9.00	12.16	15.39	18.62
Calcium (%)	0.80	0.80	0.80	0.80	0.80
Starch (%)	40.15	37.04	33.91	30.77	27.62
Fat (%)	6.21	7.47	8.72	9.97	11.23

*Grower Premix supplied the following per kg diet: Vit A 10,000,000 IU; Vit D32,000,000IU; Vit E 8,000 IU; Vit K 2,000mg; Vit B12,000 mg; Vit B25,500mg; Vit B61,200 mg; Vit B12 12 mg; Biotin30mg; Folic Acid 600 mg; Niacin 10,000 mg; Pantothenic Acid 7,000mg; Choline chloride 500,000 mg; Vit C 10,000mg; Iron 60,000 mg; Mn 80,000 mg; Cu 8,000mg; Zn 50,000 mg; Iodine 2,000 mg; Cobal 450 mg; Selenium 100 mg; Mg 100,000 mg; Anti-Oxidant 6,000 mg; PKC - Palm Kernel Cake, GNC = Groundnut cake, C.P. = Crude protein, ME = Metabolizable energy. **Ronozyme Composition of the product, sodium sulfate (52.7%), calcium carbonate (15%), kaolin (9%), dextrin and sucrose (8%), cellulose (6%) and vegetable oil (7%). Bulk density of 1,100 kg/m³. The particle size distribution of the product showed that 98% of the particles are between 150 and 1,200 µm in diameter and less than 1% of particles are below 150 µm.

Table 2. Heamatological parameters of grower pigs fed experimental diets.

Parameters	Levels of inclusion (%)					SEM (±)
	0	25	50	75	100	
	1	2	3	4	5	
PCV (%)	44.50 ^c	49.08 ^{ab}	48.45 ^{ab}	51.95 ^a	48.23 ^{ab}	2.50
RBC (×10 ⁶ /ml)	7.17 ^b	7.76 ^{ab}	7.66 ^{ab}	8.13 ^a	7.41 ^b	0.33
WBC (×10 ³ /ml)	39.50 ^a	26.45 ^d	32.75 ^b	28.20 ^{cd}	33.40 ^d	1.85
HB (g/dl)	11.48 ^b	12.90 ^a	12.85 ^a	12.98 ^a	12.60 ^d	0.54
MCV (g/dl)	59.90 ^b	63.35 ^a	63.25 ^a	64.00 ^a	63.03 ^a	1.07
MCH (pg)	15.55 ^b	16.65 ^{ab}	16.75 ^a	17.25 ^a	17.00 ^a	0.56
MCHC (%)	25.95	26.30	26.50	26.90	26.52	0.58
Neutrophils (%)	60.80	57.35	51.60	55.45	59.73	4.35

The letters a, b, c, d, e means along the same row with different superscripts are significantly ($P < 0.05$) different from each other, SEM: Standard error of mean.

Table 3. Serum biochemical parameters of grower pigs fed experimental diets.

Parameters	Levels of inclusion (%)					SEM (±)
	0	25	50	75	100	
	1	2	3	4	5	
Total protein (g/dl)	8.39 ^b	8.11 ^d	8.20 ^c	8.70 ^a	8.05 ^e	0.02
Albumin (g/dl)	3.50 ^{bc}	3.61 ^{bc}	3.40 ^{bc}	4.10 ^a	3.75 ^b	0.13
Globulin (g/dl)	4.89 ^a	4.50 ^a	4.80 ^a	4.60 ^a	4.30 ^b	0.26
Creatinine (g/dl)	1.00 ^a	0.80 ^{bc}	1.10 ^a	0.93 ^{bc}	0.87 ^{bc}	0.05
Urea (g/dl)	40.50 ^{ab}	39.50 ^a	28.50 ^c	45.50 ^a	35.50 ^b	2.90
Cholesterol (mg/dl)	118.25 ^a	116.30 ^b	114.00 ^c	110.00 ^d	109.00 ^d	0.73

The letters a, b, c, d, e means along the same row with different superscripts are significantly ($P < 0.05$) different from each other, SEM: Standard error of mean.

organisms. Hematological components are valuable in measuring the abnormality, especially with feed constituents that affect the blood as well as the physiological and health status of farm animals (Isaac et al., 2013). Results from PCV of the grower pigs were significant ($P < 0.05$) differences among levels of inclusion in the experimental diets. The result was however within the normal range for healthy grower pigs as reported by Isaac et al. (2013) that animals with good blood composition are likely to show good performance. This shows that the use of cassava peels/pkc meal (0, 25, 50, 75 and 100%) as replacement for maize had no effect on the packed cell volume of the animal blood. Packed cell volume measures the percentage composition of the blood cells in relation to other contents such as the plasma, food nutrient and is therefore useful in accessing the normal blood level in the animal.

Results obtained for RBC, WBC, Hb, MCV, MCH, and MCHC shows significant ($P < 0.05$) differences on the experimental diets. There were gradual increased from control (0) to (100%) in RBC, Hb, MCV and MCH across all levels of experimental diets. The control diet containing maize as the energy source was lower to other diets as the values were generally lower with particular references to RBC and Hb. The slight increase of hemoglobin level above that of the control from other treatments fall within the normal range and is an indication of good nutrition or adequate nutrient intake including iron, copper, vitamin or amino acid in accordance with Daramola et al. (2005), who posited that hematological values could serve as base line information for comparison of nutrient deficiency, physiology and health status of farm animals.

The serum metabolites of total protein, albumin, Globulin, creatinine, Urea and cholesterol were significantly ($P < 0.05$) affects by dietary treatments. These parameters are indicators of adequacy of protein in terms of quality and quantity in the diet. It show whether there are protein malnutrition, alterations in the dietary intake of protein and pattern of muscle utilization and possibly the extent of muscle wastage and subsequent degradation of muscle phosphorous creatinine to form creatinine according to Adeshinwa et al. (2011). The results showed that total protein, albumin and globulins in the diet supported normal protein reserves in pigs resulting efficient protein utilization (Togun et al. (2007); Unigwe et al., 2016). Serum creatinine and urea levels in animals are indicative of muscular wastage (Harper et al., 1979; Amaefule et al., 2012). Values obtained were fairly constant and comparable across the treatment groups; therefore, the animals did not suffered muscular wastage but efficient utilization of diets, ultimately resulting in high tissue deposition (Afolabi et al., 2010) across treatment groups. Cholesterol level was significantly highest ($P < 0.05$) in maize based and 25% diet, and was attributed to levels of fat deposited (Table 3). This result is in agreement with Togun et al. (2007) Unigwe et al. (2016) and Anayo (2010) that posited higher fibre diets reduced blood glucose which indicated decreased in the level of cholesterol as the level of inclusion of African yam beans (AYB) in the diet increased.

Conclusion

The hematological and serum biochemical parameters assessed fall within the normal range and is an indication of good nutrition or adequate nutrient intake. Therefore, the animals did not suffered muscular wastage but efficient utilization of diets, ultimately resulting in high tissue deposition across treatment groups. However, hematological and serum biochemical studies of cassava peels/PKC diets of grower pigs has proved that, it can be in cooperated on other monogastric animals is then suggested.

Open Access: This is an open access article distributed under the terms of the Creative Commons Attribution 4.0 License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original author(s) if the sources are credited.

REFERENCES

- Adeshinwa, A.O.K., Obi, O.O., Makanjuola, B.A., Oluwole, O.O. and Adesina, M.A. (2011). Growing pigs fed cassava peel based diet supplemented with or without farmazyme 3000 Proenx: Effect on growth, carcass and blood. *African Journal Biotechnology*, 10(14): 2791-2796, <https://www.ajol.info/index.php/ajb/article/view/137986/127556>
- Afolabi, K.D., Akinsoyini, A.O., Olajide, R. and Akinleye, S.B. (2010). Haematological parameters of the Nigerian local grower chickens fed varying dietary levels of palm kernel cake. *Proceedings of the 35th Annual Conference of the Nigeria Society for Animal Production*, pp. 247.
- Amaefule, K.U., Abasiokong, S.F., Onwudike, O.S.O. and Akinmoegun, M.B. (2012). Haematology and red blood cell osmotic stability of pigs fed graded levels of fermented cassava peel based diets. *Proceedings of the 17th Annual Conference of Animal Science Association of Nigeria (ASAN)*, pp. 152-153.
- Anayo, C. (2010). Biochemical and nutritional characteristics of African yam bean seeds (*Sphenostylis stenocarpa*) in diets for broiler chicken. Ph.D. Thesis, Department of Animal Science, Ambrose Alli University Ekpoma, Edo state, pp. 124-125.
- AOAC (2000). Association of Official Analytical Chemist. Official Methods of Analysis 17th Edition. Washington DC.
- Bamishaiye, E., Muhammed, N. and Bamishaiye, O. (2009). Haematological parameters of albino rats fed tiger nute (*Cyperus esculentus*) tuber oil meat based diets. *The internet Journal of Nutrition and Wellness*, 10(1): 58-62, <http://ispub.com/IJNW/10/1/9293>
- Daramola, J. O., Adeloye, A. A., Fatoba, T.A. and Soladoye, A.O. (2005). Haematological and serum in hematological indices in the grey breasted guinea fowl (*Numida Mealagris Gallata Pallatas*). *Nigeria Journal of Animal Production*, 18(2): 108-11.
- Etim, N.N., Williams, M.E., Enyenihi, G.E., Udo, M.D. and Offiong, E.E.A. (2013). Haematological parameters: Indicators of the

- physiological status of farm animals. *British Journal of Science*, 10(1): 33-45.
- FAO (2002). FAOSTAT statistics database. Retrieved from <http://aps.fao.org> on 5th April, 2015.
- Harper, A.E., Rodwell, V.W. and Mayers, P.A. (1979). Review of Physiological Chemistry (17th, Edition). Lang Medical California, 9422, pp. 60-81, 188-216.
- Isaac, L.J., Abah, G., Akpan, B. and Ekaette, I.U. (2013). Haematological properties of different breeds and sexes of rabbits. *Proceeding of the 18th Annual Conference of Animal Science Association of Nigeria*.
- Igene, F.U. (2006). Essentials of Pigs Production in Nigeria J.L.G Publishers, Ibadan 1st Edition.
- Khan, T.A. and Zafar, F. (2005). Haematological study in response to various doses of estrogen in broiler production. *International Journal Poultry Science*, 40(10): 748-751.
- Moseri, (2010). Interactional strategies in a multilingual context: Illustrative examples from the discourse of bargaining. Germany: Grin Verlag.
- Obi, I.U. (2002). Statistical methods of detecting differences between treatment means and research methodology issues in laboratory and field experiments, AP Express Publishing Company, Limited, pp. 13-19.
- Oboh, O.S., Igene, F.U. and Tariuwa (2004). Substitution of maize bran with cassava peel meal in the diets of growing pigs; Effect on performance and economics of production. *Proceedings of 9th Annual Conference of Animal Science Association of Nigeria*. pp. 94-96.
- Togun, V.A., Oseni, B.S.A., Ogundipe, J.A., Arewa, T.R., Homed, A.A., Ajonijebu, D.C., Oyeniran, A., Nwosisi, I. and Mustapha, F. (2007). Effects of chronic lead administration on the haematological parameters of rabbit- a preliminary study. *Proceeding of the 41st Conference of the Agriculture Society of Nigeria*, pp. 341.
- Unigwe, C.R., Marire, B.N., Omeke, B.C.O., Abonyi, F.O., Oladipo, T.A. and Adebayo, D.M. (2016). Effects of maize replaced fermented cassava peels and enzyme supplemented diet on haematology and serum biochemistry of cross bred female pigs. *International Journal of Advanced Research in Biological Sciences*, 3(6): 198-208, <https://www.ijarbs.com/pdfcopy/june2016/ijarbs27.pdf>
- Waugh, A. and Grant, A. (2018). Anatomy and physiology in health and illness. 13th Edn., Churchill, Livingstone, an Imprint of Elsevier Science, Ltd. pp. 584.